

Skyrizi IV (Risankizumab-rzaa)

PATIENT INFORMATION

Referral Status (check one): New Referral Updated Order Order Renewal

Patient Name: _____ DOB: _____

NKDA Allergies: _____ Weight _____ Please specify: lbs kg Height: _____

Patient Status (check one): New to Therapy Continuing Therapy | Last Treatment Date: _____ Next Due Date: _____

ICD-10 code (required): _____ ICD-10 description: _____

REQUIRED: Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.

PRESCRIPTION

NURSING

- TB status & date (list results here & attach clinicals)
- _____
- Baseline Liver Enzymes, including bilirubin (results)
- _____

PRE-MEDICATION ORDERS

- acetaminophen (Tylenol) 500mg | 650mg | 1000mg PO
- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- diphenhydramine (Benadryl) 25mg | 50mg | PO | IV
- methylprednisolone (Solu-Medrol) 40mg | 125mg IV
- hydrocortisone (Solu-Cortef) 100mg IV
- Other: _____
- Dose: _____ Route: _____
- Frequency: _____

Evaluate for TB prior to initiating treatment with SKYRIZI.

Hepatotoxicity in Treatment of Crohn's disease: Drug-induced liver injury during induction has been reported. Monitor liver enzymes and bilirubin levels at baseline and during induction, up to at least 12 weeks of treatment. Monitor thereafter according to routine patient management.

SPECIAL INSTRUCTIONS

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

 Provider Name (Print) Provider Signature Date